

RESEARCH ARTICLE

Impact of the North American monsoon on wildfire activity in the southwest United States

Nicholas J. Nauslar^{1,2}  | Benjamin J. Hatchett¹  | Timothy J. Brown¹ | Michael L. Kaplan¹ | John F. Mejia¹ 

¹Division of Atmospheric Sciences, Desert Research Institute, Reno, Nevada

²NOAA/NWS/NCEP Storm Prediction Center, Norman, Oklahoma

Correspondence

Nicholas J. Nauslar, NOAA/NWS/NCEP Storm Prediction Center, 120 David L. Boren Blvd., Suite 2100, Norman, OK 73072.

Email: nick.nauslar@noaa.gov

Funding information

U.S. Forest Service, Grant/Award Number: 10-CS-11130206-047

The North American monsoon (NAM) is an annual climate system phenomenon that develops over the Sierra Madre Occidental in western Mexico and spreads northwards into the southwest United States from June through September bringing large quantities of rainfall and lightning, which can vary greatly on intra- and inter-annual timescales. The timing of the NAM onset can lengthen or shorten the wildfire season in the southwest United States. Here we determine NAM onset thresholds and subsequent dates for the Southwest Area (SWA; Arizona, New Mexico, west Texas, and Oklahoma panhandle) and each SWA Predictive Services Area (PSA) April through September from 1995–2013. Various wildfire activity thresholds were defined to capture days or events associated with increased wildfire activity that are considered “busy” by wildland fire management in the context of an impact on firefighting resources. These defined thresholds allow for a unique examination of the relationship between the NAM and wildfire. Self-organizing maps (SOMs), utilizing 500-hPa geopotential heights and precipitable water, were implemented to identify atmospheric patterns contributing to the NAM onset and “busy” days for the SWA and each PSA. Map types from the SOMs analysis showed the transition to, during, and from the NAM. Northwards and eastwards displacements of the subtropical ridge over the SWA were associated with NAM onset. Restructuring of the subtropical ridge in time (i.e., amplification or breakdown) as inferred from map types over the SWA was directly associated with increased wildfire activity. By identifying atmospheric patterns pertinent to busy days of wildfire activity and a wildfire-based perspective of the NAM in the SWA, fire weather forecasters can proactively identify potential periods that may be particularly impactful on wildfire suppression resources.

KEYWORDS

dry thunderstorms, fire management, fire weather, monsoon onset, self-organizing maps, synoptic classification

1 | INTRODUCTION

The North America monsoon (NAM) affects much of North America with the largest impact in Mexico and the drylands of the southwest United States (hereafter Southwest; Figure 1a). The NAM is characterized as a large-scale atmospheric circulation that produces a distinct increase of warm-

season (June through September) precipitation over North America (Adams and Comrie, 1997; Grantz *et al.*, 2007). The NAM core region is centred over the Sierra Madre Occidental in northwestern Mexico (Douglas *et al.*, 1993; Barlow *et al.*, 1998). It can extend into the Southwest and farther north into the western United States (Figure 1a) (Bryson and Lowry, 1955; Reiter and Tang, 1984; Tang and Reiter,




FIGURE 1 (a) Map of study area and the Southwest Area outlined and shaded. (b) Study area showing the Southwest Area Predictive Service Areas (PSAs). SW01 is also known as PSA 1 [Colour figure can be viewed at wileyonlinelibrary.com]

1984; Douglas *et al.*, 1993; Higgins *et al.*, 1999; Hawkins *et al.*, 2002; Lo and Clark, 2002). The variability of NAM in the southwestern United States is large, sometimes larger than the mean warm season rainfall, and is modulated by intra-seasonal transient features including tropical easterly waves, tropical storms/cyclones, and transient upper-tropospheric troughs (Higgins *et al.*, 1998; Adams and Stensrud, 2007; Abatzoglou and Brown, 2009; Hendon *et al.*, 2011; Favors and Abatzoglou, 2012; Means, 2012; Newman and Johnson, 2012; Mejia *et al.*, 2015; Seastrand *et al.*, 2015; Pascale and Bordoni, 2016).

Higgins *et al.* (1997) describe the NAM in three phases: development, mature, and decay. The development of the NAM (May–June) is characterized by a transition of generally westerly flow associated with a boreal winter circulation shifting to a southerly and easterly flow, and moves northwards as the subtropical ridge develops over Mexico and the Southwest (Bryson and Lowry, 1955; Adams and Comrie, 1997; Grantz *et al.*, 2007). The shift in winds is in response to surface heating over the continent, especially over the elevated terrain in the NAM region (Figure 1a). As the land–ocean temperature gradient increases, air flows towards the continent increasing convergence and enhancing the mountain–valley circulation over the Sierra Madre Occidental (Douglas *et al.*, 1993; Vera *et al.*, 2006; Gochis and Higgins, 2007; Grantz *et al.*, 2007). The pronounced diurnal cycle of rainfall associated with the NAM initially develops in southern Mexico before extending north to the Sierra Madre Occidental and the Southwest (June–July) (Douglas *et al.*, 1993; Adams and Comrie, 1997; Higgins *et al.*, 1997). The heaviest precipitation occurs in July and August during the mature stage then wanes in September (Carleton *et al.*, 1990; Douglas *et al.*, 1993; Higgins *et al.*, 1997). The NAM decay phase is more gradual than onset and represents

the return to a cold season circulation with an equatorwards migration of the polar jet stream. This transition is noted by a return of the westerlies from a diurnally driven, warm-season circulation of easterly and southerly flow over the NAM region (Higgins *et al.*, 1997; Barlow *et al.*, 1998; Vera *et al.*, 2006; Cerezo-Mota *et al.*, 2011; Means, 2012).

There are several definitions for the NAM onset, but it is generally characterized by an increase of atmospheric humidity and precipitation associated with the establishment of the subtropical ridge over western Mexico and the Southwest (Adams and Comrie, 1997; Higgins *et al.*, 1997; Ellis *et al.*, 2004; Grantz *et al.*, 2007). However, a universally accepted NAM onset definition does not exist for the Southwest. NAM onset definitions include a dewpoint temperature of 54 °F at the Tucson airport (National Weather Service Weather Forecast Office Tucson, 2015), a precipitation threshold +0.5 mm/day for three consecutive days (Higgins *et al.*, 1997), or a Gulf of California sea surface temperature of 26 °C (Mitchell *et al.*, 2002).

Limited research has examined the relationship between the NAM and wildfire occurrence in the western United States (Mohrle, 2003; Brandt, 2006; Evett *et al.*, 2008). In the Southwest, wildfire occurrence increases steadily through April, but the rate accelerates in May and June before decreasing in late July through September (Brandt, 2006). The peak Southwest wildfire season occurs in June and early July due to decreasing fuel moisture, increasing temperatures, and enhanced ignition potential coinciding with drier thunderstorms before the NAM onset (Watson *et al.*, 1994; Westerling *et al.*, 2003). The NAM increases cloud-to-ground lightning, which increases “natural” wildfire ignitions, but with the corresponding increase in atmospheric and fuel moisture, wildfires often remain small as the NAM matures (Evett *et al.*, 2008; Dowdy and Mills,

2012). The increase of cloud-to-ground lightning is related to 57% of natural wildfires that occur after NAM onset (Mohrle, 2003). Moreover, Mohrle (2003) notes 55% of large wildfires (40 ha) and nearly 50% of area burned from large wildfires occur after onset demonstrating the delayed effects of NAM precipitation on fuel moisture recovery. However, wildfire occurrence abruptly decreases approximately midway through the climatological peak of lightning and precipitation (late July to early August) in Arizona and New Mexico as fuel moisture rebounds (Hall, 2007).

While a direct relationship exists between weather and wildfires, this relationship can be complex and nonlinear. Synoptic composites are often used to examine the relationships between the dominant atmospheric circulation systems and ecological factors (Hewitson and Crane, 2002). Self-organizing maps (SOMs) are utilized to identify atmospheric circulation patterns associated with different phenomena due to their propensity for objectively characterizing the complex distribution of synoptic states (Kohonen, 2001; Cavazos *et al.*, 2002; Hewitson and Crane, 2002; Reusch *et al.*, 2005; Crimmins, 2006; Reusch *et al.*, 2007; Reusch, 2011; Swales *et al.*, 2016). For example, Cavazos *et al.* (2002) showed that the NAM in southeast Arizona was dominated by three wet modes, while Crimmins (2006) identified different atmospheric patterns associated with extreme fire weather conditions in the Southwest.

Days with numerous wildfire ignitions and multiday wildfire events are of particular importance to fire meteorologists and managers due to the strain these events place on local, regional, and national suppression resources. Specifically, the Southwest Geographic Area Coordination Center, which oversees the coordination of firefighting resources in the Southwest Area (SWA) (Arizona, New Mexico, west Texas, and Oklahoma Panhandle) (Figure 1b) would bear the brunt of these impacts from these significant wildfire episodes or “busy” days as fire management describes them. Identifying atmospheric patterns pertinent to significant wildfire episodes in the SWA could provide improved prediction of significant wildfire episodes. This would facilitate more efficient allocation of firefighting resources during busy days.

In this analysis, we examine various definitions and thresholds for determining the NAM onset and significant wildfire episodes (busy days) within the SWA and its individual Predictive Services Areas (PSAs) (Figure 1b). We suggest that SOMs can help examine the intricate nonlinear relationships between the NAM and critical fire weather patterns leading to significant wildfire episodes. We implement SOMs to identify synoptic patterns that correspond to the NAM onset and significant wildfire episodes including those atmospheric patterns that are prevalent before, during, and after these events. SOMs help infer the associated physical climate processes between the NAM and wildfire activity.

2 | DATA

Data were collected for the SWA spanning April through September from 1995 to 2013 unless otherwise specified. Southwest Predictive Services defined the PSAs as having similar historical wildfire occurrences, fuel types, fuel indices, and climates (i.e., temperature and precipitation characteristics) (Figure 1b). They also determined large fire thresholds based on historical fire records and suppression needs (Table 1).

We used daily, 4-km surface gridded data of precipitation, minimum and maximum relative humidity (RH, RHmin, RHmax), specific humidity (q), minimum and maximum temperature (T_{min} , T_{max}), and wind speed from the University of Idaho METDATA (Abatzoglou, 2013). Vapour pressure deficit (VPD) was defined as the difference between saturated vapour pressure and ambient vapour pressure. Saturated vapour pressure (e_s ; in mb) was calculated using Tetens’s (1930) formula as follows:

$$e_s = 6.11 * 10^{\left(\frac{7.5 * T}{237.3 + T}\right)},$$

where T is temperature in °C. Vapour pressure deficit (VPD; in mb) was then calculated as

$$VPD = e_s * RH / 100,$$

where RH is relative humidity in percent. A single daily PSA value for each variable was calculated by averaging all grid point values of each variable within a PSA.

Daily 700 and 500 hPa winds, geopotential heights, and q and daily precipitable water (PW) were obtained from the National Center for Environmental Prediction (NCEP) North American Regional Reanalysis (NARR; Mesinger *et al.*, 2006) and downloaded from the Earth System Research Laboratory Physical Sciences Division (<https://www.esrl>).

TABLE 1 Large wildfire threshold for each PSA in the SWA as determined by southwest PS

PSA	Large wildfire threshold (acres)
SW01	100
SW02	300
SW03	300
SW04	100
SW05	50
SW06S	100
SW06N	500
SW07	100
SW08	300
SW09	300
SW10	100
SW11	100
SW12	100
SW13	2000
SW14	200
SW14N	2000

noaa.gov/psd/data/gridded/data.narr.html). Despite some known limitations of NARR (Dominguez *et al.*, 2008; Becker *et al.*, 2009; Cerezo-Mota *et al.*, 2011), numerous studies have shown NARR representative when examining synoptic circulation and moisture across the study region (e.g., Dominguez *et al.*, 2008; Dominguez and Kumar, 2008; Turrent and Cavazos, 2012; Doubler *et al.*, 2015; Radhakrishna *et al.*, 2015). We used a subset of daily NARR data that spanned from 85°W to 131°W longitude and 18°N to 49.5°N latitude.

Wildfire data were obtained from the Fire Program Analysis (FPA) quality controlled wildfire database (Short, 2015), which has the size, location, discovery date, ignition cause (i.e., type) among other information for each wildfire. Wildfires were categorized by type (all, lightning, human) and size (all, large) for each PSA, and large wildfire criteria were established from large wildfire thresholds determined by Southwest Predictive Services (Table 1). The Program for Climate, Ecosystem, and Fire Applications at the Desert Research Institute provided proprietary National Lightning Detection Network[®] CG lightning data.

3 | METHODS

3.1 | The North American monsoon onset

The predictors selected for determining the NAM onset were the daily averaged PSA values of q , VPD, cloud-to-ground lightning strikes (LS), and precipitation. These variables were chosen because each represents atmospheric moisture or thunderstorm coverage, which are hallmarks of the NAM onset. To eliminate potential “pre-monsoonal” moisture surges, time periods of 3–10 days and variable thresholds that exceeded the 22 year, April through September PSA mean or median values for q and precipitation were considered. Periods where VPD values were less than their 22-year mean or median were also considered. A threshold of 100 LS was chosen due to the Storm Prediction Center using it as its forecast lightning density threshold and after examining various thresholds between 25 and 200 LS. The primary technique for identifying NAM onset used six consecutive days of greater (less) than or equal to the 22-year median PSA value of q (VPD) and six consecutive days of at least 100 lightning strikes occurring concurrently (onset 1). Other combinations of variables were evaluated that substituted precipitation for q (onset 2) or VPD (onset 3). The NAM onset date for each PSA was determined by taking the first instance that met the onset 1, onset 2, or onset 3 requirements after May 31. We constrained the onset after May 31 to minimize the influences of spring storms and also pre-monsoon moisture surges. While noting the subjectivity of these methods, they do represent the fundamental weather and climate shift the NAM exerts on the SWA and produces

realistic results based on operational fire weather forecaster experience. For each PSA, the 10th percentile, 90th percentile, median, inter-quartile range, and ± 1 median absolute deviation onset dates were calculated.

3.2 | Significant wildfire episodes

The FPA wildfire occurrence data were organized by PSA and separated into categories based on size and type. If a day exceeded the 95th (99th) percentile for number of wildfires and large wildfires or at least two (three) large wildfires occurred for a PSA, it was considered a “busy day” and exceeded the Busy95_PSA (Busy99_PSA) threshold. If a day exceeded the 95th (99th) percentile for number of large wildfires for the SWA, it was considered a “busy day” and exceeded the Busy95_SW (Busy99_SW) threshold. Periods of 2, 3, 7, and 10 days that exceeded the 95th and 99th percentiles for large wildfires for the SWA over those respective time periods were identified as “busy events.” These thresholds and periods represent a level of fire activity that can begin to stress local and regional wildfire suppression resources. The busy days and events were examined primarily to determine (a) when these episodes occurred most frequently and (b) to examine synoptic patterns during significant wildfire episodes (i.e., busy days or events).




FIGURE 2 Self-organizing map node distances for the 4×5 grid 4,000 iteration solution. The blue hexagons represent the nodes and the red lines connect neighbouring nodes. The colours in the regions containing the red lines indicate the distances between nodes with the darker colours representing larger distances and lighter colours representing smaller distances [Colour figure can be viewed at wileyonlinelibrary.com]

3.3 | Self-organizing maps

As noted in the introduction, SOMs are a nonparametric analysis technique based on neural networks that extract generalized patterns from gridded data defined over a region (Kohonen, 2001). SOMs can process multivariate, multidimensional input on a user-defined grid (e.g., 4×3 ; 12 nodes) by creating a spatially organized set of generalized patterns of variability from the input data (Reusch *et al.*, 2005; 2007; Reusch, 2011). SOMs capture the most notable atmospheric circulation patterns. Because the analysis is performed over daily time increments, such patterns include those modes related to synoptic scale variations (Cavazos *et al.*, 2002; Reusch *et al.*, 2007). This process is iterative

and continues until the node locations and densities converge on a solution and become stable (i.e., stop changing) (Hewitson and Crane, 2002). The iterative SOMs process clusters nodes in regions where there is more information while spanning the distribution of data and placing similar (dissimilar) nodes closer to (farther from) each other (Hewitson and Crane, 2002). The SOMs divide a continuum of atmospheric patterns into a small number of categories, which are spatially organized by similarity with similar patterns closer to each other and dissimilar patterns further away from each other on the user-defined grid (Hewitson and Crane, 2002; Reusch, 2011). By identifying atmospheric circulation patterns, SOMs allows a frequency analysis of




FIGURE 3 Twenty SOM reference nodes (MTs) in the 4×5 SOM grid arrangement for 500 hPa geopotential heights (m), contoured and colour shaded every 30 m [Colour figure can be viewed at wileyonlinelibrary.com]

each pattern, preferred transitions, and the ability to associate patterns with events (Cavazos *et al.*, 2002; Reusch *et al.*, 2005; Reusch *et al.*, 2007).

Although we did initially examine results using principal component analysis and *k*-means clustering, the SOMs methodology was ultimately chosen because it has been shown to be more robust, accurate, and able to provide better discriminating power of variability patterns in large, nonlinear data sets (Reusch *et al.*, 2005; 2007). Furthermore, SOMs allows more discrimination among subtle variations of similar atmospheric patterns (Hewitson and Crane, 2002). Kohonen (2001), Hewitson and Crane (2002), (Reusch *et al.*, 2005; 2007), and Reusch (2011) provide further detailed explanations of

SOMs and the implementation of SOMs for meteorological and climatological applications.

SOMs were implemented using various packages from the Neural Network Toolbox in MATLAB® (Mathworks, 2017). The input data consisted of standardized daily values of NARR 500-hPa geopotential heights, 700-hPa geopotential heights, and PW on a 175×120 grid (85° – 131° W; 18° – 49.5° N) for 183 days (April 1 to September 30) over 22 years (1992–2013). SOMs require standardized data when using multiple variables with varying scales (Reusch *et al.*, 2007; Reusch, 2011). All input data were included together for training because the primary goal of SOMs is to use generalized patterns determined from the input data as recommended by Reusch (2011). Asymmetrical grid




FIGURE 4 Twenty SOM reference nodes (MTs) in the 4×5 SOM grid arrangement for precipitable water (mm), contoured and colour shaded every 5 mm [Colour figure can be viewed at wileyonlinelibrary.com]

configurations of nodes are recommended (Kohonen, 2001; Reusch *et al.*, 2007) thus node grid arrays of 4×3 (rows by columns), 5×3 , and 4×5 were examined at 500, 1,000, 2,000, and 4,000 iterations or until there was little variation of node densities and locations (i.e., stable). Sammon maps of multidimensional data on the different SOMs grids (not shown) and matching the input data to the final reference nodes of these grids displayed how dissimilar the nodes were and the distances among those nodes (Figure 2) (Sammon, 1969; Reusch *et al.*, 2007). Plots of neighbour node distances and examining the dates associated with each node were also helpful in examining the SOMs solutions (Figure 2). Each node was assigned a number (1–20) via SOMs, which helped facilitate analysis of the solutions. After a stable solution was reached, each day was classified with a node as a best match, which was composited into a map type (MT) for a particular meteorological variable (Reusch, 2011). After each day was classified, a frequency analysis was applied to all days, busy days/events, and onset days where applicable to determine if individual MTs or MT transitions were more associated with certain types of events.

4 | RESULTS

4.1 | Atmospheric map types and NAM “life cycle”

The Sammon maps and neighbour node distances and densities for the three examined node grid configurations were all generally acceptable. Each grid configuration satisfied recommended SOMs mathematical requirements and

resolved expected atmospheric patterns for the study area during the examined period (e.g., Hewitson and Crane, 2002). However, the 4×5 grid resolved two patterns corresponding to busy and NAM onset days that were not well represented by the 4×3 and 5×3 grids. The 20 node configuration versus the 12 or 15 node configuration helped discriminate relatively subtle variations in the atmospheric circulation patterns especially at the intersection of troughs and ridges, which helped identify these two additional important MTs.

A stable solution was reached after 2,000 iterations on the 4×5 SOMs grid, but the 4,000 iteration version was utilized due to the slightly better coverage and shape from the Sammon map depiction (not shown). Figures 3 and 4 display the resulting 20 composite MTs from the SOMs analysis. Map type 1 (MT 1, e.g., Figure 3, MT 1) for 500-hPa geopotential heights directly corresponds to MT 1 for both 700-hPa geopotential heights (not shown; similar to 500 hPa patterns) and PW (e.g., Figure 4, MT 1) because all three variables were included in the same SOMs analysis. Anomaly MTs were created, but as they were very similar in appearance, they are not shown here.

Each MT represents a node from the SOMs solution using the 4×5 grid with 4,000 iterations. Figure 2 displays the distances (e.g., magnitude of dissimilarity) among adjacent nodes, and each one of those nodes corresponds directly to the MTs in Figures 3 and 4. For example, the node at the very top, left of Figure 2, directly corresponds to MT 1. The number assigned to each node increases left to right across the columns and top to bottom across the rows. This layout helps compare the nodes/MTs mathematically and meteorologically.

TABLE 2 Map type (MT) occurrence by month summed for the period of 1995–2013 and associated median lightning strike (LS) total decile

	April	May	June	July	August	September	Median LS decile
MT 1	134	1	0	0	0	0	1
MT 2	104	18	0	0	0	0	1
MT 3	98	43	0	0	0	0	1
MT 4	54	64	2	0	0	0	3
MT 5	103	36	0	0	0	0	1
MT 6	107	29	0	0	0	0	1
MT 7	35	137	15	0	0	3	2
MT 8	1	93	79	0	0	6	4
MT 9	16	124	33	0	0	13	4
MT 10	8	106	51	0	0	9	2
MT 11	0	7	164	19	8	30	4
MT 12	0	4	76	17	27	83	5
MT 13	0	4	88	12	7	126	4
MT 14	0	0	29	55	70	76	5
MT 15	0	0	34	83	86	36	6
MT 16	0	0	34	146	97	8	8
MT 17	0	16	29	1	0	160	3
MT 18	0	0	14	52	110	78	6
MT 19	0	0	8	170	145	19	7

Ascending numbered MTs follow a general progression of increasing geopotential heights and PW (Figures 3 and 4, MTs 1–20). Tighter geopotential height gradients, deeper troughs with lower heights, and drier conditions are illustrated at lowered numbered MTs (Figures 3 and 4, MTs 1–10) over the SWA and are most frequent in April and May (Table 2). Ascending numbered MTs generally occur in roughly chronological order through August. This trend parallels the amplification of subtropical ridge and increased moisture over northwestern Mexico and the SWA, which is indicative of the NAM (Figures 3 and 4 and Table 2) (e.g., Douglas *et al.*, 1993; Adams and Comrie, 1997; Stensrud *et al.*, 1997). As the NAM decays in September, lower numbered MTs become more frequent indicating a decrease in geopotential heights and drier conditions (Figures 3 and 4, MTs 12–13).

April is characterized by deeper troughs and dry conditions prevailing across the SWA (Table 2 and Figures 3 and 4, MTs 1–2, 5–6). The most prevalent MTs in May depict various longitudinal extents of the subtropical ridge

axis that extends from the south and east over Mexico and the Gulf of Mexico (Table 2 and Figure 3, MTs 7–10). Drier conditions remain over most of the SWA, but PW increases on the western and eastern coasts of Mexico (Figure 4, MTs 7–10). Geopotential heights continue to increase and the variation between troughs and ridges over the SWA maximizes during June (Table 2 and Figure 3, MTs 8, 11–13). In June, the subtropical ridge axis is centred over western and northwestern Mexico and PW increases in extent and magnitude across Mexico, the Plains, and the SWA (Figures 3 and 4, MTs 11–13; e.g., PSAs 13°, 14 N, 14°S). The increase in lower-tropospheric moisture from the Gulf of Mexico is due in large part to the Low-Level Jet (Stensrud, 1996; Higgins *et al.*, 1997; Stensrud *et al.*, 1997). During July, the transition to a ridge-dominated atmospheric pattern is complete with the establishment of the Four Corners High (i.e., MT 20). This pattern continues through August with easterlies shifting northwards bringing ample moisture across much of Mexico and into the SWA (Table 2 and Figures 3 and




FIGURE 5 PSA median onset date for the three different onset metrics from 1995 to 2013 [Colour figure can be viewed at wileyonlinelibrary.com]

TABLE 3 Median, inter-quartile range (IQR; number of days), median absolute deviation (MAD; number of days), 10th percentile onset date, and 90th percentile onset day-of-year (DOY) by PSA for the three onset metrics using specific humidity (q), vapour pressure deficit (VPD), precipitation (precip), and lightning strikes (LS) from 1995 to 2013

	SW01	SW02	SW03	SW04	SW05	SW06S	SW06N	SW07	SW08	SW09	SW10	SW11	SW12	SW13	SW14S	SW14N
Onset 1 (q , VPD, LS)	Jul 24	Jul 14	Jul 30	Jul 20	Jul 18	Jul 8	Jul 18	Jul 15	Jul 11	Jul 6	Jul 13	Jul 8	Jul 12	Jun 30	Jun 26	Jul 27
IQR	35	25	32	19	19	10	19	21	15	17	29	25	26	21	34	52
MAD	17	10	14	10	10	6	11	11	8	7	13	13	14	11	11	19
90th	Aug 21	Aug 11	Aug 30	Aug 12	Aug 7	Jul 20	Aug 3	Aug 3	Jul 24	Jul 30	Jul 28	Aug 1	Aug 4	Jul 23	Aug 4	Aug 20
10th	Jul 4	Jul 3	Jul 4	Jul 4	Jul 3	Jun 29	Jun 30	Jul 1	Jun 28	Jun 20	Jun 16	Jun 11	Jun 18	Jun 4	Jun 14	Jun 5
Onset 2 (VPD, Precip, LS)	Jul 24	Jul 14	Jul 30	Jul 20	Jul 18	Jul 9	Jul 18	Jul 15	Jul 11	Jul 6	Jul 17	Jul 13	Jul 14	Jul 4	Jul 2	Jul 2
IQR	35	25	32	19	19	10	17	25	16	17	28	20	36	34	39	53
MAD	17	10	14	10	10	6	5	12	8	7	14	10	18	19	16	26
90th	Aug 21	Aug 11	Aug 30	Aug 12	Aug 7	Jul 20	Aug 3	Aug 9	Jul 25	Jul 30	Aug 3	Aug 3	Aug 12	Aug 9	Aug 14	Aug 10
10th	Jul 4	Jul 3	Jul 4	Jul 4	Jul 3	Jun 29	Jun 30	Jul 1	Jun 28	Jun 12	Jun 25	Jun 14	Jun 14	Jun 5	Jun 14	Jun 2
Onset 3 (q , Precip, LS)	Jul 18	Jul 11	Jul 13	Jul 13	Jul 8	Jul 2	Jul 8	Jul 5	Jul 1	Jul 2	Jun 29	Jul 6	Jul 8	Jun 15	Jun 22	Jul 6
IQR	20	16	20	18	15	10	15	16	14	11	21	18	32	22	40	37
MAD	12	9	9	9	7	7	8	8	8	7	6	9	13	8	15	20
90th	Aug 7	Jul 31	Aug 16	Jul 31	Jul 17	Jul 15	Jul 25	Jul 14	Jul 15	Jul 24	Jul 19	Jul 18	Aug 10	Jul 11	Aug 1	Aug 6

4, MTs 15–16, 18–20). In September, the subtropical ridge becomes less amplified with MTs demonstrating weaker ridges and troughs over the Southwest (Figure 3, MTs 12–13, 17).

The four most frequent MTs (Figure 3, e.g., MTs 16, 18–20) show the SWA under the influence of the subtropical ridge, which is expected during the NAM (Adams and Comrie, 1997; Higgins *et al.*, 1997). PW increases in magnitude and northwards extent during ridging confirming prior research (Figures 3 and 4, e.g., MTs 15–16, 18–20) (Carleton *et al.*, 1990; Higgins *et al.*, 1999; Cerezo-Mota *et al.*, 2011). Lightning strikes also increase during these ridging patterns. MTs 16, 20 have the highest median lightning decile (Table 2) and MTs 16, 19–20 are most associated with days exceeding the 70th percentile of LS across the SWA.

4.2 | Quantifying NAM onset by Predictive Services Area

Figure 5 and Table 3 provide the median onset date, inter-quartile range, median absolute deviation, 10th percentile onset date, and 90th percentile onset date by PSA for the three onset metrics implemented. Onset1 (q , VPD, and lightning) and onset 2 (VPD, precipitation, and lightning) generally produce the same results with some variation across eastern New Mexico and west Texas (Figure 5). Onset 3 (q , precipitation, and lightning) uniformly shows earlier median onset dates for every PSA except PSA SW14N (Figure 5), and onset 3 is less temporally variable than onset 1 and onset 2 (Table 3). Most onset dates for all the metrics occur in early to mid-July with a few instances of late June onset dates in some of the eastern New Mexico and west Texas PSAs (Figure 5). Depending on the PSA and metrics, the inter-quartile range can range from 10 to 53 days although most inter-quartile ranges total less than 35 days (Table 3). The median absolute deviations for onset are generally between 1 and 2 weeks with a few exceptions totalling around 3 weeks. The 10th and 90th percentile onset dates are generally separated by three to six weeks (Table 3), which indicates that while NAM onset varies inter-annually, it typically begins between late June and early August. Adjacent PSAs tend to have onset dates within a week of each other and occasionally on the same day.

Ridging, including a northwards displacement of the subtropical ridge, southerly and easterly flow (inferred from assuming geostrophic flow at 500 hPa), increased moisture, and LS are associated with onset days (Figures 3 and 4, MTs 15–16, 18–20 and Table 4). MTs 15–16, 18–20 total 28.7% of all days examined, but occur on 75.0% of onset days (Table 4). MTs 1–7 are never associated with onset for a PSA. These MTs generally depict lower heights and a trough at 500 hPa with less LS and drier conditions over the SWA. In these cases, moisture advection is limited from the eastern Pacific due to cooler waters and numerous mountain ranges

TABLE 4 Showing the percentage of occurrence for each map type (MT) for all days, Busy95_SW, Busy99_SW, and onset 1 days and the median lightning strikes (LS) decile (Dec) for each MT

	Count	All days (%)	Busy95_SW (%)	Busy99_SW (%)	Onset1	Median LS decile
MT 1	135	3.4	0.5	0.0	0.0	1
MT 2	122	3.0	2.3	3.5	0.0	1
MT 3	141	3.5	2.7	1.8	0.0	1
MT 4	120	3.0	6.4	8.8	0.0	3
MT 5	139	3.5	2.3	1.8	0.0	1
MT 6	136	3.4	2.7	3.5	0.0	1
MT 7	190	4.7	4.1	3.5	0.0	2
MT 8	179	4.4	5.5	5.3	1.8	4
MT 9	186	4.6	3.2	3.5	0.7	4
MT 10	174	4.3	2.7	1.8	0.0	2
MT 11	228	5.7	11.0	14.0	2.8	4
MT 12	207	5.1	8.2	10.5	3.9	5
MT 13	237	5.9	2.7	5.3	8.5	4
MT 14	230	5.7	2.7	5.3	6.0	5
MT 15	239	5.9	5.0	1.8	9.9	6
MT 16	285	7.1	12.8	7.0	19.0	8
MT 17	206	5.1	2.3	0.0	1.4	3
MT 18	254	6.3	3.2	0.0	12.3	6
MT 19	342	8.5	9.6	10.5	18.7	7

TABLE 5 Mean number of wildfires per year for the SWA by type and size for 1992–2013

Total wildfires per year	4,064
Total large wildfires per year	108
Lightning caused wildfires per year	1,972
Lightning caused large wildfires per year	62
Human caused wildfires per year	2,092

between the Pacific and the SWA (Table 4 and Figures 3 and 4, MTs 1–7). These MTs also predominantly occur in April and May and rarely in June through August (Table 4).

4.3 | Wildfire occurrence patterns and associated map types

Annually, 51.5% of wildfires are human caused, but 57.2% of all large wildfires are lightning caused (Table 5). Wildfires increase in frequency into July before declining, confirming prior results from Brandt (2006). Seasonally, 3.5% of wildfires ignited in April through June become large, while only 1.9% of wildfires become large in July through September (Table 6). This finding is consistent with results

from Mohrle (2003), who showed 57% of natural wildfires, but only 37% of total area burned from natural wildfires occurred after onset (early to mid-July). Human-related wildfires drop after onset also confirming findings from Mohrle (2003).

More than half of the days that exceeded the Busy95_PSA or Busy99_PSA threshold occurred before July 1, and more than 82% of days that exceeded these thresholds occurred before the respective PSA's annual NAM onset 1 date (Table 7). When examining the entire Southwest, most of the busy days occur before July 1 and almost all of them occur before August (Table 7). PSAs SW06S, SW08, and SW12 frequently had large fires when the Busy95_SW and Busy99_SW thresholds were exceeded (Table 8). On average, four PSAs had at least one large fire when exceeding the Busy99_SW threshold indicating widespread wildfire activity across the SWA during the busiest events.

MTs 4, 11–12, 16, 19–20 are most associated with days exceeding either the Busy95_SW or Busy99_SW threshold (Table 4). MTs 4, 11–12 are more associated with days exceeding either busy threshold than onset 1 days, while the converse is true for MTs 16, 19–20 (Table 4). Crimmins

TABLE 6 Total SWA wildfires by type and month for 1992–2013

	Wildfires	Large wildfires	Lightning-caused wildfires	Lightning-caused large wildfires	Human wildfires	Human-caused large wildfires
April	8,318	300	744	67	7,574	233
May	14,789	417	3,348	151	11,441	266
June	21,443	822	8,774	476	12,669	346
July	25,687	496	17,367	375	8,320	121
August	13,899	267	10,545	229	3,354	38

TABLE 7 Displaying the number (count) of occurrences for each type of event from 1992 to 2013. Also shows the percentage of events that transpired before onset (if applicable) and before July 1. Shows number of occurrences for each type of event by month

	Count	% before onset	% before 1 July	April	May	June	July	August	September
Busy95_PSA	298	85.23	62.4	28	40	118	76	33	3
Busy99_PSA	292	82.53	55.1	25	31	105	98	30	3
Busy95_SW	219		66.2	23	31	91	51	22	1
Busy99_SW	57		77.2	6	10	28	11	2	0
Event_2_95	108		70.4	13	19	44	24	8	0
Event_3_95	83		72.3	8	17	35	17	6	0
Event_7_95	49		73.5	6	7	23	11	2	0
Event_10_95	36		77.8	3	8	17	7	1	0
Event_2_99	27		81.5	2	5	15	5	0	0
Event_3_99	17		88.2	1	4	10	2	0	0
Event_7_99	11		81.8	0	2	7	2	0	0
Event_10_99	10		80.0	0	2	6	2	0	0

(2006) noted three strong southwesterly flow patterns that are associated with extreme fire weather conditions in the Southwest, and those patterns are captured by our SOMs. Specifically, MT 11 is similar to node 10 from Crimmins (2006) and is associated with 11.0 and 14.0% of Busy95_SW and Busy99_SW days, respectively (Table 4). Several MTs 1, 17–18 are rarely associated (<1%) with either busy threshold and those three MTs are distinct compared to each other (Figure 3 and Table 4).

5 | DISCUSSION

Busy days and events across the SWA are generally associated with latitudinally suppressed upper-tropospheric ridging (subtropical ridge axis in western/northwestern Mexico) (e.g., MTs 4, 8, 12), southwest flow with higher 500-hPa geopotential heights (e.g., MT 11), or latitudinally amplified ridging (subtropical ridge axis in SWA) (e.g., MTs 16, 19–20). The suppressed ridge and southwest flow with higher geopotential heights MTs (Figure 3, MTs 4, 8, 12) have PW values of 10–25 mm across most of the SWA, but deeper moisture is confined farther south in Mexico (Figure 4, MTs 4, 8, 12). Due to the complex terrain in the SWA and northern Mexico, mid-tropospheric (700–400 hPa) moisture often arrives before lower-tropospheric moisture. Additionally, MTs 4, 8, 11–12 depict atmospheric flow that would be less conducive for lower-tropospheric moisture advection from the Gulf of California into the SWA compared to MTs 16, 19–20 (Figures 3,4 and 6). Strong diurnal heating and upslope flow can create a deeply mixed boundary layer and thermally induced circulations that can utilize increased mid-tropospheric moisture and produce thunderstorms (Hales, 1977). With lower-tropospheric moisture confined farther south, an environment conducive for dry thunderstorms develops with characteristic inverted-v soundings where a dry sub-cloud layer and high LCL forms an inverted-v between the dewpoint

and temperature profiles (Figures 3,4 and 6, MTs 4, 8, 11–12 and Table 4) (Hall, 2007; Wallmann *et al.*, 2010; Nauslar *et al.*, 2013). Transient upper-tropospheric short-wave troughs and easterly waves also act to increase buoyancy and augment coverage of thunderstorms by lowering heights and temperatures aloft.

MTs with amplified ridging over the SWA also have higher PW values and more LS across the SWA (Figures 3 and 4, MTs 15–16, 18–20 and Table 2). As the ridge builds northwards over the SWA, easterly and southerly flow transports moisture from the Gulf of Mexico across northern Mexico and into the SWA. Additionally, southerly flow increases lower-tropospheric moisture advection from the Gulf of California into the SWA. The increased moisture creates a more unstable atmosphere and helps trigger thunderstorms beyond the typical terrain-favoured locations (e.g., Mogollon Rim and Southern Rockies) (Gebremichael *et al.*, 2007; Gochis and Higgins, 2007; Nesbitt *et al.*, 2008). MTs 15–16, 18–20 that are characterized by amplified ridging are most associated with onset and often occur in July and August when the NAM typically begins and matures. However, MTs 16, 19–20 also often correspond to busy days (Figures 3 and 4). This could be due to the increased amount and spatial coverage of lightning (Table 4), which enhance ignition potential across a broader area. Enhanced lightning also likely increases fire activity across multiple PSAs given the presence of receptive fuels. Both outcomes strain wildfire suppression resources. Furthermore, MTs 16, 19–20 are more likely to help produce busy wildfire days and events if they occur earlier in the season (e.g., before onset). At this time, peak fuel dryness could offset the increase of atmospheric moisture to help produce the most active part of the wildfire season. The term “Firesoon” has been used to describe these early moisture intrusions due to their propensity for triggering thunderstorms that ignite wildfires (Brown, 2002). Not coincidentally, 3.8% of wildfires in June become large while only 1.9% of wildfires in July become large (Table 6). Additionally, 75.6% of busy events occur

before July 1 and 83.9% of all busy days occur before onset (Table 7). Wildfires, especially large wildfires, that begin well before onset strain wildfire suppression resources because they are likely to continue growing until the NAM becomes established.

While associating MTs with busy days and events is important, the atmospheric patterns that precede and follow those days are equally important. Each MT represents a climatology of similar atmospheric patterns that typically occur from April through September across the SWA. By analysing MTs assigned to specific days around busy days and events, observations and inferences about the progression of atmospheric patterns (e.g., MTs) can be gleaned. Four general atmospheric pattern progressions emerge when examining MTs around busy days and events for the SWA: (a) zonal or southwest flow transition to ridging; (b) variation between suppressed and amplified ridging over the SWA; (c) zonal or southwest flow transitioning to ridging followed by a return to zonal or southwest flow; and (d) ridging over the SWA with the ridge axis exhibiting east–west movement (Figure 7).

The MT progressions around busy days and events usually occur on timescales of days to weeks. This underscores the importance of progressive wave patterns characterized by alternating upper-tropospheric troughs and ridges. MTs with upper-tropospheric troughing and zonal or southwesterly flow over the SWA have lower PW values and tighter height gradients across the SWA. These conditions translate to dry and windy surface conditions that help cure fuels and promote a receptive fuel bed for wildfire ignition and spread. The MTs characterized by upper-tropospheric ridging and higher 500-hPa geopotential heights are associated with higher LS and PW values across the SWA. Higher 500-hPa geopotential heights are also indicative of warmer surface temperatures. This variation between periods of hot, dry, and windy surface conditions and increased moisture and lightning appears to drive much of the fire weather and occurrence across the SWA. Specifically, MT progressions that represent a transition from upper-tropospheric ridging to troughing are reminiscent of the breakdown of the upper-tropospheric ridge, a known critical fire weather pattern (Werth *et al.*, 2011). The breakdown involves decreasing geopotential heights with an eastwards or equatorwards shift of the ridge axis that can lead to the following pattern change: (a) hot and dry surface conditions; (b) increasing atmospheric instability and moisture leading to thunderstorms; and (c) dry and windy surface conditions. That progression is conducive for large wildfire occurrence as lightning occurs with receptive fuels, which is followed by dry and windy conditions that promote fire spread. The periodicity of these patterns varies, and future work could entail examining the duration and frequency of these patterns more closely.

TABLE 8 Displays the number of times a PSA had at least one large wildfire when the Busy99_PSA or Busy95_PSA threshold was exceeded. Also shows the percentage representing the proportion for each PSA when having at least one large wildfire when the Busy99_SW or Busy95_SW threshold was exceeded

	SW01	SW02	SW03	SW04	SW05	SW06S	SW06N	SW07	SW08	SW09	SW10	SW11	SW12	SW13	SW14S	SW14N
Busy99_SW	7 12.3%	20 35.1%	4 7.0%	6 10.5%	14 24.6%	39 68.4%	15 26.3%	6 10.5%	23 40.4%	12 21.1%	8 14.0%	9 15.8%	28 49.1%	11 19.3%	13 22.8%	13 22.8%
Busy95_SW	24 11.0%	46 21.0%	14 6.4%	16 7.3%	42 19.2%	120 54.8%	47 21.5%	22 10.0%	70 32.0%	25 11.4%	27 12.3%	18 8.2%	61 27.9%	31 14.2%	38 17.4%	31 14.2%




FIGURE 6 700 hPa geopotential heights contoured every 20 m (black, solid lines) and specific humidity colour shaded every 1 g/kg for MTs 4, 8, 11–12 [Colour figure can be viewed at wileyonlinelibrary.com]

Holdovers are wildfires that are ignited but remain small (i.e., single tree) until favourable atmospheric conditions develop and subsequently promote fire growth. A wildfire's discovery date and its ignition date may not match, and holdovers are difficult to quantify and predict (Anderson, 2002). Holdovers and large wildfire growth after the discovery date introduce errors into our analysis. However, utilizing the busy day and event thresholds mitigate some of these issues because these consider relatively rare events (<5% of occurrence) and deemphasize total acres burned. Future work could use the Monitoring Trends in Burn Severity database to calculate area burned on a daily timescale and apply the same rigorous atmospheric pattern and fuel indices analysis. However, such an analysis would be confined to examining only larger wildfires due to the spatial resolution and accuracy of this database (Kolden *et al.*, 2012).

The methods implemented in this analysis produce realistic and meaningful annual NAM onset dates by PSA. This helps frame wildfire occurrence in the SWA. Generally, onset progresses from east to west and south to north across the SWA, but with large inter-annual variability. The differences in the onset dates between Arizona and western New Mexico from eastern New Mexico and west Texas could be attributed to a precipitation dipole that exists between the Plains (central United States) and the mountainous Southwest (Figure 5) (Higgins *et al.*, 1997; Higgins *et al.*, 1999; Mitchell *et al.*, 2002; Adams and Stensrud, 2007; Stensrud, 2013). MTs 15–16, 18–20 are most associated with the NAM onset and depict southerly flow over most of the

SWA and a northwards shift of easterlies over northern Mexico leading to increased lightning and PW across the SWA (Figures 3 and 4, MTs 15–16, 18–20) (Table 4). These MTs are very similar to the SOM nodes that Cavazos *et al.* (2002) found to be most associated with wet NAM modes for southeast Arizona (or “bursts,” as in Carleton (1986) and Watson *et al.* (1994)). MT 17 is curiously unrelated to NAM onset, but 500-hPa heights are lower than other MTs associated with onset, and the ridge axis is shifted west limiting moisture advection from the Gulf of California. This induces drier west–northwest flow over most of the Southwest, which confirms prior research (Table 4) (Carleton, 1986; Carleton *et al.*, 1990; Higgins *et al.*, 1999; Cerezo-Mota *et al.*, 2011).

6 | CONCLUSIONS

Using atmospheric reanalysis data, gridded surface weather observations, and historical wildfire data sets we demonstrated that the most frequently occurring synoptic patterns before and during the NAM modulate large wildfire occurrence. Several MTs or MT progressions are associated with significant wildfire episodes (i.e., busy days or events) (Figure 7):

- Zonal or southwest flow transition to ridging.
- Fluctuation between suppressed and amplified ridging over the SWA.




FIGURE 7 Most frequent MT progressions associated with increased wildfire activity (contours have same values as Figure 3) [Colour figure can be viewed at wileyonlinelibrary.com]

- Zonal or southwest flow transitioning into ridging followed by a return to zonal or southwest flow.
- Variation between suppressed and amplified ridging over the SWA with the ridge axis exhibiting east–west movement.

If these patterns develop before the NAM onset, and thus coincide with peak fuel and atmospheric dryness, then the likelihood of significant wildfire episodes increases. A

method for determining the NAM onset by PSA as directly related to wildfire was also established corresponding to observed, and to some extent predictable, atmospheric patterns that promote increased moisture and lightning. We provided novel results documenting MTs directly associated with significant wildfire episodes and NAM onset in the SWA.

The results of this work are intended to provide decision support information and improve understanding of

atmospheric processes associated with NAM and their impact on wildfire activity. Improved understanding of this association may benefit operational fire meteorologists and managers by providing identifiable atmospheric patterns associated with increased wildfire activity that are particularly impactful on suppression resources and can lead to overall improvements in planning and logistical strategies (Ray *et al.*, 2007).

ACKNOWLEDGEMENTS

We would like to thank the USDA Forest Service for providing the funding (10-CS-11130206-047), and Vaisala for allowing the use of lightning data for wildfire applications. We would also like to thank Chuck Maxwell, John Saltenberger, and Terry Marsha for their input and advice for the paper. Finally, we would like to thank David Reusch and Michael Crimmins for their expertise and help on SOMs, John Abatzoglou for helping obtain and organize the MET-DATA, and Andrew Joros for helping obtain and organize the NARR data set. We would also like to thank the two anonymous reviewers for their thorough and constructive comments, which greatly improved the manuscript.

ORCID

Nicholas J. Nauslar  <https://orcid.org/0000-0003-1600-6311>

Benjamin J. Hatchett  <https://orcid.org/0000-0003-1066-3601>

John F. Mejia  <https://orcid.org/0000-0001-6727-5541>

REFERENCES

- Abatzoglou, J.T. (2013) Development of gridded surface meteorological data for ecological applications and modelling. *International Journal of Climatology*, 33, 121–131.
- Abatzoglou, J.T. and Brown, T.J. (2009) Influence of the Madden–Julian Oscillation on summertime cloud-to-ground lightning activity over the continental United States. *Monthly Weather Review*, 137, 3596–3601.
- Adams, D.K. and Comrie, A.C. (1997) The North American monsoon. *Bulletin of the American Meteorological Society*, 78, 2197–2213.
- Adams, J.L. and Stensrud, D.J. (2007) Impact of tropical easterly waves on the North American monsoon. *Journal of Climate*, 20, 1219–1238.
- Anderson, K. (2002) A model to predict lightning-caused fire occurrences. *International Journal of Wildland Fire*, 11, 163–172.
- Barlow, M., Nigam, S. and Berbery, E.H. (1998) Evolution of the North American monsoon system. *Journal of Climate*, 11, 2238–2257.
- Becker, E.J., Berbery, E.H. and Higgins, R.W. (2009) Understanding the characteristics of daily precipitation over the United States using the North American regional reanalysis. *Journal of Climate*, 22, 6268–6286.
- Brandt, R.R. (2006) *The North American monsoon system in southern Arizona*. Dissertation, Department of Geography and Regional Development, University of Arizona, 130 pp.
- Brown, T.J. (2002) The North American “Firesoon.” Report on research opportunities for climate and society interactions in the North American monsoon region. In: Tucson, A.Z., Ray, A.J. and Webb, R.S. (Eds.) *Workshop on Applications and Human Dimensions of Monsoon Research, June 18–20, 2001*. Tucson, AZ: NOAA OAR, 64 pp.
- Bryson, R.A. and Lowry, W.P. (1955) Synoptic climatology of the Arizona summer precipitation singularity. *Bulletin of the American Meteorological Society*, 36, 329–339.
- Carleton, A.M. (1986) Synoptic-dynamic character of “bursts” and “breaks” in the southwest U.S. summer precipitation singularity. *Journal of Climatology*, 6, 605–623.
- Carleton, A.M., Carpenter, D.A. and Weser, P.J. (1990) Mechanisms of interannual variability of the southwest United States summer rainfall maximum. *Journal of Climate*, 3, 999–1015.
- Cavazos, T., Comrie, A.C. and Liverman, D.M. (2002) Intraseasonal variability associated with wet monsoons in southeast Arizona. *Journal of Climate*, 15, 2477–2490.
- Cerezo-Mota, R., Allen, M. and Jones, R. (2011) Mechanisms controlling precipitation in the northern portion of the North American monsoon. *Journal of Climate*, 24, 2771–2783.
- Crimmins, M.A. (2006) Synoptic climatology of extreme fire-weather conditions across the southwest United States. *International Journal of Climatology*, 26, 1001–1016.
- Dominguez, F. and Kumar, P. (2008) Precipitation recycling variability and eco-climatological stability—a study using NARR data. Part I: central U.S. plains ecoregion. *Journal of Climate*, 21, 5165–5186.
- Dominguez, F., Kumar, P. and Vivoni, E.R. (2008) Precipitation recycling variability and eco-climatological stability—a study using NARR data. Part II: North American monsoon region. *Journal of Climate*, 21, 5187–5203.
- Doubler, D.L., Winkler, J.A., Bian, X., Walters, C.K. and Zhong, S. (2015) An NARR-derived climatology of southerly and northerly low-level jets over North America and coastal environments. *Journal of Applied Meteorology and Climatology*, 54, 1596–1619.
- Douglas, M.W., Maddox, R.A., Howard, K. and Reyes, S. (1993) The Mexican monsoon. *Journal of Climate*, 6, 1665–1677.
- Dowdy, A.J. and Mills, G.A. (2012) Atmospheric and fuel moisture characteristics associated with lightning-attributed fires. *Journal of Applied Meteorology and Climatology*, 51, 2025–2037.
- Ellis, A.W., Saffell, E.M. and Hawkins, T.W. (2004) A method for defining monsoon onset and demise in the southwestern USA. *International Journal of Climatology*, 24, 247–265.
- Evert, R.R., Mohrle, C.R., Hall, B.L., Brown, T.J. and Stephens, S.L. (2008) The effect of monsoonal atmospheric moisture on lightning wildfire ignitions in southwestern North America. *Agricultural and Forest Meteorology*, 148, 1478–1487.
- Favors, J.E. and Abatzoglou, J.T. (2012) Regional surges of monsoonal moisture into the southwestern United States. *Monthly Weather Review*, 141, 182–191.
- Gebremichael, M., Vivoni, E.R., Watts, C.J. and Rodríguez, J.C. (2007) Submesoscale spatiotemporal variability of North American monsoon rainfall over complex terrain. *Journal of Climate*, 20, 1751–1773.
- Gochis, D.J. and Higgins, R.W. (2007) The path to improving predictions of the North American monsoon. *US CLIVAR Variations*, 5.
- Grant, K., Rajagopalan, B., Clark, M. and Zagona, E. (2007) Seasonal shifts in the North American monsoon. *Journal of Climate*, 20, 1923–1935.
- Hales, J.E. (1977) On the relationship of convective cooling to nocturnal thunderstorms at Phoenix. *Monthly Weather Review*, 105, 1609–1613.
- Hall, B.L. (2007) Precipitation associated with lightning-ignited wildfires in Arizona and New Mexico. *International Journal of Wildland Fire*, 16, 242–254.
- Hawkins, T.W., Ellis, A.W., Skindlov, J.A. and Reigle, D. (2002) Intra-annual analysis of the North American snow cover–monsoon teleconnection: seasonal forecasting utility. *Journal of Climate*, 15, 1743–1753.
- Hendon, H.H., Sperber, K.R., Waliser, D.E. and Wheeler, M.C. (2011) Modeling monsoon intraseasonal variability: from theory to operational forecasting. *Bulletin of the American Meteorological Society*, 92, ES32–ES35.
- Hewitson, B.C. and Crane, R.G. (2002) Self-organizing maps: applications to synoptic climatology. *Climate Research*, 22, 13–26.
- Higgins, R.W., Yao, Y. and Wang, X.L. (1997) Influence of the North American monsoon system on the U.S. summer precipitation regime. *Journal of Climate*, 10, 2600–2622.
- Higgins, R.W., Mo, K.C. and Yao, Y. (1998) Interannual variability of the U.S. summer precipitation regime with emphasis on the southwestern monsoon. *Journal of Climate*, 11, 2582–2606.
- Higgins, R.W., Chen, Y. and Douglas, A.V. (1999) Interannual variability of the North American warm season precipitation regime. *Journal of Climate*, 12, 653–680.
- Kohonen, T. (2001) *Self-Organizing Maps*. Berlin Heidelberg: Springer, 503 pp.
- Kolden, C.A., Lutz, J.A., Key, C.H., Kane, J.T. and van Wageningen, J.W. (2012) Mapped versus actual burned area within wildfire perimeters: characterizing the unburned. *Forest Ecology and Management*, 286, 38–47.

- Lo, F. and Clark, M.P. (2002) Relationships between spring snow mass and summer precipitation in the southwestern United States associated with the North American monsoon system. *Journal of Climate*, 15, 1378–1385.
- Mathworks. (2017) Self-organizing maps. Available at: <https://www.mathworks.com/help/nnet/self-organizing-maps.html> [Accessed 15th September 2017].
- Means, J.D. (2012) GPS precipitable water as a diagnostic of the North American monsoon in California and Nevada. *Journal of Climate*, 26, 1432–1444.
- Mejia, J.F., Douglas, M.W. and Lamb, P.J. (2015) Observational investigation of relationships between moisture surges and mesoscale to large-scale convection during the North American monsoon. *International Journal of Climatology*, 36, 2555–2569.
- Mesinger, F., DiMego, G., Kalnay, E., Mitchell, K., Shafran, P.C., Ebisuzaki, W., Jović, D., Woollen, J., Rogers, E., Berbery, E.H., Ek, M.B., Fan, Y., Grumbine, R., Higgins, W., Li, H., Lin, Y., Manikin, G., Parrish, D. and Shi, W. (2006) North American regional reanalysis. *Bulletin of the American Meteorological Society*, 87, 343–360.
- Mitchell, D.L., Ivanova, D., Rabin, R., Brown, T.J. and Redmond, K. (2002) Gulf of California sea surface temperatures and the North American monsoon: mechanistic implications from observations. *Journal of Climate*, 15, 2261–2281.
- Mohrle, C.R. (2003) *The southwest monsoon and relation to fire occurrence*. MS thesis, Department of Physics/Division Of Atmospheric Sciences (DRI), University of Nevada, Reno, 97 pp. Available at: <http://www.cefa.dri.edu/Publications/charliethesis.pdf> [Accessed 15th June 2018].
- National Weather Service Weather Forecast Office Tucson. (2015) Monsoon information index. Available at: http://www.wrh.noaa.gov/twc/monsoon/monsoon_info.php [Accessed 12th November 2015].
- Nauslar, N.J., Kaplan, M.L., Wallmann, J. and Brown, T.J. (2013) A forecast procedure for dry thunderstorms. *Journal of Operational Meteorology*, 1, 200–214.
- Nesbitt, S.W., Gochis, D.J. and Lang, T.J. (2008) The diurnal cycle of clouds and precipitation along the Sierra Madre Occidental observed during NAME-2004: implications for warm season precipitation estimation in complex terrain. *Journal of Hydrometeorology*, 9, 728–743.
- Newman, A. and Johnson, R.H. (2012) Mechanisms for precipitation enhancement in a North American monsoon upper-tropospheric trough. *Journal of the Atmospheric Sciences*, 69, 1775–1792.
- Pascale, S. and Bordoni, S. (2016) Tropical and extratropical controls of Gulf of California surges and summertime precipitation over the southwestern United States. *Monthly Weather Review*, 144, 2695–2718. <https://doi.org/10.1175/MWR-D-15-0429.1>.
- Radhakrishna, B., Fabry, F., Braun, J.J. and Van Hove, T. (2015) Precipitable water from GPS over the continental United States: diurnal cycle, intercomparisons with NARR, and link with convective initiation. *Journal of Climate*, 28, 2584–2599.
- Ray, A.J., Garfin, G.M., Wilder, M., Vásquez-León, M., Lenart, M. and Comrie, A.C. (2007) Applications of monsoon research: opportunities to inform decision making and reduce regional vulnerability. *Journal of Climate*, 20, 1608–1627.
- Reiter, E.R. and Tang, M. (1984) Plateau effects on diurnal circulation patterns. *Monthly Weather Review*, 112, 638–651.
- Reusch, D.B. (2011) Nonlinear climatology and paleoclimatology: capturing patterns of variability and change with self-organizing maps. *Physics and Chemistry of the Earth*, 35, 329–340.
- Reusch, D.B., Alley, R.B. and Hewitson, B.C. (2005) Relative performance of self-organizing maps and principal component analysis in pattern extraction from synthetic climatological data. *Polar Geography*, 29, 227–251.
- Reusch, D.B., Alle, R.B. and Hewitson, B.C. (2007) North Atlantic climate variability from a self-organizing map perspective. *Journal of Geophysical Research*, 112, D02104.
- Sammon, J.W. (1969) A nonlinear mapping for data structure analysis. *IEEE Transactions on Computers*, 18, 401–409.
- Seastrand, S., Serra, Y., Castro, C. and Ritchie, E. (2015) The dominant synoptic-scale modes of North American monsoon precipitation. *International Journal of Climatology*, 35, 2019–2032.
- Short, K.C. (2015) *Spatial Wildfire Occurrence Data for the United States, 1992–2013 [FPA_FOD_20150323]*, 3rd edition. Fort Collins, CO: Forest Service Research Data Archive.
- Stensrud, S.J. (1996) Importance of low-level jets to climate: a review. *Journal of Climate*, 9, 1698–1711. [https://doi.org/10.1175/15200442\(1996\)009<1698:IOLLJT.2.0.CO;2](https://doi.org/10.1175/15200442(1996)009<1698:IOLLJT.2.0.CO;2).
- Stensrud, D.J. (2013) Upscale effects of deep convection during the North American monsoon. *Journal of the Atmospheric Sciences*, 70, 2681–2695.
- Stensrud, D.J., Gall, R.L. and Nordquist, M.K. (1997) Surges over the Gulf of California during the Mexican monsoon. *Monthly Weather Review*, 125, 417–437.
- Swales, D., Alexander, M. and Hughes, M. (2016) Examining moisture pathways and extreme precipitation in the U.S. Intermountain West using self-organizing maps. *Geophysical Research Letters*, 43, 1727–1735.
- Tang, M. and Reiter, E.R. (1984) Plateau monsoons of the Northern Hemisphere: a comparison between North America and Tibet. *Monthly Weather Review*, 112, 617–637.
- Tetens, O. (1930) Über einige meteorologische Begriffe. *Zeitschrift für Geophysik*, 6, 297–309.
- Turrent, C. and Cavazos, T. (2012) A numerical investigation of wet and dry onset modes in the North American monsoon core region. Part I: A regional mechanism for interannual variability. *Journal of Climate*, 25, 3953–3969.
- Vera, C., et al. (2006) Toward a unified view of the American monsoon systems. *Journal of Climate*, 19, 4977–5000.
- Wallmann, J., Milne, R., Smallcomb, C. and Mehle, M. (2010) Using the June 21, 2008 California lightning outbreak to improve dry lightning forecast procedures. *Weather and Forecasting*, 25, 1447–1462.
- Watson, A.I., Holle, R.L. and López, R.E. (1994) Cloud-to-ground lightning and upper-air patterns during bursts and breaks in the southwest monsoon. *Monthly Weather Review*, 122, 1726–1739.
- Werth, P.A., Potter, B.E., Clements, C.B., Finney, M.A., Goodrick, S.L., Alexander, M.E., Cruz, M.G., Forthofer, J.A. and McAllister, S.S. (2011) *Synthesis of knowledge of extreme fire behavior: volume I for fire managers*. Portland, OR: US Department of Agriculture, Forest Service, Pacific Northwest Research Station. General technical report number: PNW-GTR-854.
- Westerling, A.L., Brown, T.J., Gershunov, A., Cayan, D.R. and Dettinger, M.D. (2003) Climate and wildfire in the Western United States. *Bulletin of the American Meteorological Society*, 84(5), 595–604.

How to cite this article: Nauslar NJ, Hatchett BJ, Brown TJ, Kaplan ML, Mejia JF. Impact of the North American monsoon on wildfire activity in the southwestern United States. *Int J Climatol*. 2019;39: 1539–1554. <https://doi.org/10.1002/joc.5899>